

## Arcegel Matrix High Concentration, Phenol Red-Free, LDEV-Free

### Product description

Arcegel Matrix High Concentration, Phenol Red-Free, LDEV-Free is a soluble basement membrane preparation extracted from EHS mouse tumors rich in extracellular matrix proteins. Its main components are laminin, type IV collagen, heparan sulfate proteoglycan (HSPG), nestin as well as growth factors such as TGF-beta, EGF, IGF, FGF, tissue plasminogen activator and other growth factors contained in EHS tumors. At room temperature, it aggregates to form a biologically active three-dimensional matrix, which simulates the structure, composition, physical properties and functions of the cell basement membrane in vivo, which is beneficial to the culture and differentiation of cells in vitro. It can be used for studies of cell morphology, biochemical function, migration, invasion and gene expression.

Arcegel Matrix is a sterile product with a concentration of more than 18 mg/mL, which meets a variety of experimental requirements, including in vivo tumorigenesis, angiogenesis studies, tumor cell migration, 3D cell studies and other applications.

### Specifications

Catalog Number	C231006E/C231006S
Specifications	5 mL/10 mL

### Properties

Properties	Parameters
Product Line	Arcegel
Product specifications	5/10 mL
Classification	High Concentration
Product Type	Basement Membrane Matrix
Form	Frozen
Species	EHS Mouse Tumors
Concentration	≥18 mg/mL
Endotoxin Level	Low
Phenol Red Indicator	None
Serum Level	None
LDEV Detection	None

## Storage

Transported on dry ice. Stored at -20°C with a shelf life of 2 years.

## Notes

1. After thawing, gently shake the reagent bottle to evenly disperse the Arcegel matrix gel.
2. All procedures involving aliquoting, usage, etc., must be conducted in a sterile environment. The bottle cap can be wiped with 70% ethanol and left to air dry.
3. Experimental equipment (e.g., pipette tips, product tubes) in contact with the product must be pre-cooled to ensure that the Arcegel matrix gel remains uniform.
4. Cells can grow on the surface of the Arcegel matrix gel at a thickness of 0.5 mm or within the three-dimensional matrix of the Arcegel matrix gel at a thickness of 1 mm. Over-diluted Arcegel matrix gel will form a non-gelatinous protein layer, suitable for cell adhesion but not for cell differentiation studies.
5. Arcegel quickly gels at temperatures between 22~35°C. The gelled Arcegel matrix gel can return to a liquid state after 24~48 hours at 4°C.
6. After melting, aliquot the Arcegel into multiple small tubes. All aliquots must be placed in pre-cooled cryovials, rapidly frozen, and stored to avoid repeated thawing and freezing.
7. For your safety and health, please wear lab coats and disposable gloves when handling.
8. For research use only.

## Instructions

### 1. Thawing and preservation of Arcegel basement membrane/Matrix

**【Note】** Arcegel Matrix is very sensitive to temperature and must not be frozen and thawed repeatedly. The dispensing of Arcegel Matrix and the preparation before gelation must be performed on ice (4°C), because a slight increase of temperature may cause gelation, resulting in uneven Matrix or affecting subsequent gelation. Tubes or pipette tips used for holding must be pre-cooled.

- 1) After receiving the product, if you do not use it temporarily, please directly store the whole bottle at -20°C (do not store it in a frost-free refrigerator).
- 2) For the first use, put the entire bottle of Arcegel Matrix in an ice box and put it at 4°C overnight to fully melt.

### 2. The use of Arcegel basement membrane/Matrix

Matrix gels rapidly at 22~35°C. In order to ensure the gel-forming performance and stability of Arcegel Matrix, the final dilution concentration should not be lower than 3 mg/mL (the

concentration of Arcegel Matrix liquid varies from batch to batch).

Arcegel Matrix can be diluted in serum-free medium and should be used immediately after dilution.

- 1) Thin gel preparation method
  - a. After melting, mix the Arcegel Matrix with a pre-cooled pipette tip.
  - b. Put the culture plate to be used on ice, and add Arcegel Matrix at a concentration of 50  $\mu$  L/cm<sup>2</sup> growth area.
  - c. Incubate at 37°C for 30 min, then the plate can be used.
- 2) Thick gel preparation method
  - a. After melting, mix the Arcegel Matrix with a pre-cooled pipette tip.
  - b. Put the culture plate to be used on ice, mix the cultured cells with Arcegel Matrix, and use a cold pipette tip to suspend the cells evenly. Arcegel Matrix was added at a concentration of 150~200  $\mu$ L/cm<sup>2</sup> growth area.
  - c. Incubate at 37°C for 30 min and then the cell culture medium can be added. Cells can also grow on top of this thick gel.
- 3) Thin-layer coating method
  - a. After melting, mix the Arcegel Matrix with a pre-cooled pipette tip.
  - b. Dilute Arcegel Matrix to the desired concentration with serum-free medium. It is recommended to do a gradient experiment according to the specific experiment to determine the optimal coating concentration.
  - c. Add the diluted Arcegel Matrix to the culture vessel to be coated, and the coating amount covers at least all growth surfaces of the cells. Incubate for 1 hour at room temperature.
  - d. Remove uncoagulated and bound Arcegel Matrix and rinse gently with serum-free medium. The plate is now ready for use.

**【Note】** Arcegel Matrix-coated plates are best used on the same day, but can also be adjusted according to specific applications. After adding the medium, the coated plates can be stored at 37°C for up to 1 week.