

Recombinant Human TARC/CCL17 Protein

Product Information

Product Name	Cat#	Size
	90963ES08	5 µg
Recombinant Human TARC/CCL17 Protein	90963ES60	100 µg
	90963ES76	500 µg

Product Description

CCL17, also known as thymus and activation-related chemokine (TARC), a small cytokine belonging to the C-C chemokine family. Among CC chemokine family members, CCL17 has approximately 24 - 29% amino acid sequence identity with RANTES, MIP-1 alpha, MIP-1 beta, MCP-1, MCP-2, MCP-3 and I-309. CCL17 is constitutively expressed in thymus, and at a lower level in lung, colon, and small intestine. CCL17 is also transiently expressed in phytohemagglutinin-stimulated peripheral blood mononuclear cells. CCL17 has been shown to be chemotactic for T cell lines but not monocytes or neutrophils. CCL17 was identified to be a specific functional ligand for CCR4, a receptor that is selectively expressed on T cells.

Product Properties

Synonyms	MGC138273, SCYA17, SCYA17MGC138271
Accession	Q92583
GeneID	6361
Source	E.coli-derived Human TARC/CCL17, Ala24-Ser93.
Molecular Weight	Approximately 8.1 kDa.
AA Sequence	ARGTNVGREC CLEYFKGAIP LRKLKTWYQT SEDCSRDAIV FVTVQGRAIC SDPNNKRVKN AVKYLQSLER S
Tag	None
Physical Appearance	Sterile Filtered White lyophilized (freeze-dried) powder.
Purity	> 97% by SDS-PAGE and HPLC analyses.
Biological Activity	The biological activity determined by a chemotaxis bioassay using human T-lymphocytes is in a concentration range of 1.0-10 ng/mL. Fully biologically active when compared to standard.
Endotoxin	< 1.0 EU per 1µg of the protein by the LAL method.
Formulation	Lyophilized from a 0.2 µm filtered concentrated solution in 20 mM PB, pH 7.4, 150 mM NaCl. We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom.
Reconstitution	Reconstitute in sterile distilled water or aqueous buffer containing 0.1% BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at ≤ -20°C. Further dilutions should be made in appropriate buffered solutions.

Shipping and Storage

The products are shipped with ice pack and can be stored at -20°C for 1 year.

1 month, 2 to 8 °C under sterile conditions after reconstitution.

3 months, -20 °C under sterile conditions after reconstitution.

Recommend to aliquot the protein into smaller quantities when first used and avoid repeated freeze-thaw cycles.

Cautions

1. Avoid repeated freeze-thaw cycles.
2. For your safety and health, please wear lab coats and disposable gloves for operation.
3. For research use only.