

Recombinant Human Beta-defensin 3 (Human BD-3)

Product Information

Product Name	Cat#	Size
	92503ES08	5 µg
Recombinant Human Beta-defensin 3 (Human BD-3)	92503ES60	100 µg
	92503ES76	500 µg

Product Description

Defensins (alpha and beta) are cationic peptides with antimicrobial activity against Gram-negative and Gram-positive bacteria, fungi, and enveloped viruses. They are 2-6 kDa proteins and take important roles in innate immune system. On the basis of their size and pattern of disulfide bonding, mammalian defensins are classified into alpha, beta and theta categories. β -Defensins contain a six-cysteine motif that forms three intra-molecular disulfide bonds. Four human β -defensins have been identified and they are expressed on some leukocytes and at epithelial surfaces. Because β -defensins is cationic peptides, they can therefore interact with the membrane of invading microbes, which are negative due to lipopolysaccharides (LPS) and lipoteichoic acid (LTA) found in the cell membrane. Especially, they have higher affinity to the binding site compared to Ca^{2+} and Mg^{2+} ions. Furthermore, they can affect the stability of the membrane.

Product Properties

Synonyms	BD-3 Protein, DEFB-3 Protein, DEFB103 Protein, DEFB3 Protein, BD-3 Protein, BD3 Protein, BP-3 Protein, BP3 Protein
Accession	P81534
GeneID	55894
Source	E.coli-derived human Beta-defensin 3 protein, Gly23-Lys67
Molecular Weight	Approximately 5.2 kDa.
AA Sequence	GIINTLQKYY CRVRGGRC AV LSCLPK EEQI GKCSTRGRKC CRRKK
Tag	None
Physical Appearance	Sterile Filtered White lyophilized (freeze-dried) powder.
Purity	> 98 % by SDS-PAGE and HPLC analyses.
Biological Activity	Fully biologically active when compared to standard. The ED_{50} as determined by anti-microbial activity against E.coli. is less than 30 µg/ml, corresponding to a specific activity of > 33.3 IU/mg.
Endotoxin	< 1.0 EU per 1µg of the protein by the LAL method.
Formulation	Lyophilized from a 0.2 µm filtered concentrated solution in 20 mM PB, pH 7.4, 130 mM NaCl. We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom.
Reconstitution	Reconstitute in sterile distilled water or aqueous buffer containing 0.1% BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at $\leq -20^{\circ}\text{C}$. Further dilutions should be made in appropriate buffered solutions.

Shipping and Storage

The products are shipped with ice pack and can be stored at -20°C to -80°C for 1 year.

Recommend to aliquot the protein into smaller quantities when first used and avoid repeated freeze-thaw cycles.

Cautions

1. Avoid repeated freeze-thaw cycles.
2. For your safety and health, please wear lab coats and disposable gloves for operation.
3. For research use only!