

Recombinant Mouse Interleukin-33 (Mouse IL-33)

Product Information

Product Name	Cat#	Size
	90157ES10	10 µg
Recombinant Mouse Interleukin-33 (Mouse IL-33)	90157ES60	100 µg
	90157ES76	500 µg

Product Description

IL-33, also known as NF-HEV and DVS 27, is a 17.5 kDa proinflammatory protein that may also regulate gene transcription. DVS 27 was identified as a gene that is up-regulated in vasospastic cerebral arteries. NF-HEV was described as a nuclear factor that is preferentially expressed in the endothelial cells of high endothelial venules relative to endothelial cells from other tissues. IL-33 was identified based on sequence and structural homology with IL-1 family cytokines. DVS 27, NF-HEV, and IL-33 share 100% amino acid sequence identity. IL-33 is constitutively expressed in smooth muscle and airway epithelia. It is up-regulated in arterial smooth muscle, dermal fibroblasts, and keratinocytes following IL-1 alpha or IL-1 beta stimulation. Similar to IL-1, IL-33 can be cleaved in vitro by caspase-1, generating an N-terminal fragment that is slightly shorter than the C-terminal fragment. The N-terminal portion of full length IL-33 contains a predicted bipartite nuclear localization sequence and a homeodomain-like helix-turn-helix DNA binding domain. By immunofluorescence, full length IL-33 localizes to the nucleus in HUVECs and transfectants. The C-terminal fragment, corresponding to mature IL-33, binds and triggers signaling through mast cell IL-1 R4/ST2L, a longtime orphan receptor involved in the augmentation of Th2 cell responses. A ternary signaling complex is formed by the subsequent association of IL-33 and ST2L with IL-1 RAcP. Stimulation of Th2 polarized lymphocytes with mature IL-33 in vitro induces IL-5 and IL-13 secretion. In vivo administration of mature IL-33 promotes increased production of IL-5, IL-13, IgE, and IgA, as well as splenomegaly and inflammatory infiltration of mucosal tissues. Full length and mature mouse IL-33 share approximately 55% and 90% aa sequence identity with human and rat IL-33, respectively. Mouse IL-33 shares less than 25% aa sequence identity with other IL-1 family proteins.

Product Properties

Synonyms	IL-1F11, NF-HEV
Accession	Q8BVZ5
GeneID	77125
Source	E.coli-derived Mouse IL-33, Ser109-Ile266.
Molecular Weight	Approximately 17.5 kDa.
AA Sequence	SIQGTSLLTQ SPASLSTYND QSVSFVLENG CYVINVDDSG KDQEQQVLL RYYESPCPAS QSGDGVGK LMVNMSPIKD TDIWLHANDK DYSVELQRGD VSPPEQAFFV LHKKSSDFVS FECKNLPGTY IGVKDNQLAL VEEKDESCNN IMFKLSKI
Tag	None
Physical Appearance	Sterile Filtered White lyophilized (freeze-dried) powder.
Purity	> 98% by SDS-PAGE and HPLC analyses.
Biological Activity	The ED ₅₀ as determined by a cell proliferation assay using murine D10S cells is less than 0.5 ng/mL, corresponding to a specific activity of > 2.0 × 10 ⁶ IU/mg. Fully biologically active when compared to standard.
Endotoxin	< 1.0 EU per 1µg of the protein by the LAL method.

Formulation	Lyophilized from a 0.2 μ m filtered solution in PBS, and 1 mM EDTA. We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom.
Reconstitution	Reconstitute in sterile distilled water or aqueous buffer containing 0.1% BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at $\leq -20^{\circ}\text{C}$. Further dilutions should be made in appropriate buffered solutions.

Shipping and Storage

The products are shipped with ice pack and can be stored at -20°C to -80°C for 1 year.

Recommend to aliquot the protein into smaller quantities when first used and avoid repeated freeze-thaw cycles.

Cautions

1. Avoid repeated freeze-thaw cycles.
2. For your safety and health, please wear lab coats and disposable gloves for operation.
3. For research use only!