

# Recombinant Mouse Mucosae-associated Epithelial

## Chemokine/CCL28 (Mouse MEC/CCL28)

#### **Product Information**

Product Name	Cat#	Size
Recombinant Mouse Mucosae-associated Epithelial Chemokine/CCL28 (Mouse MEC/CCL28)	90994ES08	5 μg
	90994ES60	100 μg
	90994ES76	500 μg

## **Product Description**

MEC is a secreted CC chemokine expressed primarily by epithelial cells of the bronchioles, salivary gland, mammary gland and colon. MEC signals through the CCR10 receptor, and chemoattracts resting CD4, CD8 T-cells and eosinophils. MEC contains six cysteines, including the four highly conserved cysteine residues present in CC chemokines. Recombinant Murine MEC is a 12.6 kDa protein containing 111 amino acid residues.

## **Product Properties**

Synonyms	CCK1		
Accession	Q9JIL2		
GeneID	56838		
Source	E.coli-derived Mouse CCL28 protein, Ser20-Arg130.		
Molecular Weight	Approximately 12.6 kDa		
AA Sequence	SEAILPMASS CCTEVSHHVS GRLLERVSSC SIQRADGDCD LAAVILHVKR RRICISPHNR		
	TLKQWMRASE VKKNGRENVC SGKKQPSRKD RKGHTTRKHR TRGTHRHEAS R		
Tag	None		
Physical Appearance	Sterile Filtered White lyophilized (freeze-dried) powder.		
Purity	> 97% by SDS-PAGE and HPLC analyses.		
Biological Activity	Fully biologically active when compared to standard. The biological activity determined by a chemotaxis		
	bioassay using murine lymphocytes is in a concentration range of 1.0-10.0 ng/ml.		
Endotoxin	<1 EU/μg of protein as determined by LAL method.		
Formulation	Lyophilized from a 0.2 µm filtered concentrated solution in 20 mM PB, pH 7.4, 150 mM NaCl.		
Reconstitution	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom.		
	Reconstitute in sterile distilled water or aqueous buffer containing 0.1% BSA to a concentration of 0.1-1.0		
	mg/mL. Stock solutions should be apportioned into working aliquots and stored at $\leq$ -20°C. Further		
	dilutions should be made in appropriate buffered solutions.		

## **Shipping and Storage**

The products are shipped with ice pack and can be stored at  $-20^{\circ}$ C to  $-80^{\circ}$ C for 1 year.

Recommend to aliquot the protein into smaller quantities when first used and avoid repeated freeze-thaw cycles.

#### **Cautions**

1. Avoid repeated freeze-thaw cycles.

www.yeasen.com Page 1 of 2



- 2. For your safety and health, please wear lab coats and disposable gloves for operation.
- 3. For research use only.

www.yeasen.com Page 2 of 2